

The Journal of Threatened Taxa (JoTT) is dedicated to building evidence for conservation globally by publishing peer-reviewed articles online every month at a reasonably rapid rate at www.threatenedtaxa.org. All articles published in JoTT are registered under [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/) unless otherwise mentioned. JoTT allows unrestricted use, reproduction, and distribution of articles in any medium by providing adequate credit to the author(s) and the source of publication.

Journal of Threatened Taxa

Building evidence for conservation globally

www.threatenedtaxa.org

ISSN 0974-7907 (Online) | ISSN 0974-7893 (Print)

NOTE

RECORD OF A 10-YEAR OLD EUROPEAN WILDCAT *FELIS SILVESTRIS SILVESTRIS* SCHREBER, 1777 (MAMMALIA: CARNIVORA: FELIDAE) FROM MT. ETNA, SICILY, ITALY

Stefano Anile, Sebastien Devillard, Clayton Kent Nielsen & Mario Lo Valvo

17 February 2020 | Vol. 12 | No. 2 | Pages: 15272–15275

DOI: 10.11609/jott.5484.12.2.15272-15275



SMALL WILD CATS SPECIAL ISSUE



For Focus, Scope, Aims, Policies, and Guidelines visit <https://threatenedtaxa.org/index.php/JoTT/about/editorialPolicies#custom-0>

For Article Submission Guidelines, visit <https://threatenedtaxa.org/index.php/JoTT/about/submissions#onlineSubmissions>

For Policies against Scientific Misconduct, visit <https://threatenedtaxa.org/index.php/JoTT/about/editorialPolicies#custom-2>

For reprints, contact <ravi@threatenedtaxa.org>

The opinions expressed by the authors do not reflect the views of the Journal of Threatened Taxa, Wildlife Information Liaison Development Society, Zoo Outreach Organization, or any of the partners. The journal, the publisher, the host, and the partners are not responsible for the accuracy of the political boundaries shown in the maps by the authors.

Member



Publisher & Host





Record of a 10-year old European Wildcat *Felis silvestris silvestris* Schreber, 1777 (Mammalia: Carnivora: Felidae) from Mt. Etna, Sicily, Italy

Stefano Anile¹ , Sebastien Devillard² , Clayton Kent Nielsen³ & Mario Lo Valvo⁴

^{1,3} Cooperative Wildlife Research Laboratory, University of Southern Illinois, Carbondale 62901, USA.

² Université Claude Bernard Lyon 1, CNRS, Laboratoire de Biométrie et Biologie Evolutive, F-69100, Villeurbanne, France.

⁴ Laboratorio di Zoologia applicata, Dipartimento STEBICEF, University of Palermo, 90123, Palermo, Italy.

¹ stefanoanile@yahoo.it (corresponding author), ² sebastien.devillard@univ-lyon1.fr, ³ kezo92@siu.edu, ⁴ mario.lovalvo@unipa.it

Information on longevity is key to the understanding of population biology of a species (Healy et al. 2014) and is being collected by researchers across taxa (Cutler 1979; Wilkinson & South 2002; De Magalhaes & Costa 2009; Gonzalez-Lagos et al. 2010; Tidiere et al. 2016). Longevity data of wildlife can be collected through long-term monitoring programs (Smith et al. 2017), but is often difficult to apply to rare and elusive species, such as small carnivores. Alternatively, longevity data can also be collected both from dead animals by using cementum annuli to estimate the age of death (Kamler & Macdonald 2006) and from species held in captivity, although it is well-known that captive animals generally live longer than those in the wild (Ricklefs & Cadena 2007; Tidiere et al. 2016).

Within the Felidae there is a consistent bias in the scientific community to study larger species rather than smaller ones (Brodie 2009; Macdonald et al. 2010; Anile & Devillard 2015, 2018), and hence scientific data on life history traits for smaller species are lacking. Given the aforementioned constraints, knowledge of longevity of small carnivores in the wild is rare. To our knowledge, the only longevity study of European Wildcats is that of

Hartmann (2005) in Switzerland, where captive animals attained 12–16 years of age. Here we report our recent finding of a European Wildcat recaptured by camera traps after nine years on Mt. Etna in Sicily, Italy.

Our study area was located on Mt. Etna in Sicily, Italy (Fig. 1A), the highest active volcano in Europe and a World Heritage Site by UNESCO in 2013 (UNESCO World Heritage Center 2019). Extensive descriptions of our study area are reported elsewhere (Anile et al. 2014, 2019).

We have been using camera traps to survey the Wildcat population on Mt. Etna since 2006 (Anile et al. 2009, 2010, 2012a,b, 2014, 2019). Extensive details about the methods and materials used in the surveys related to the present study are reported in Anile et al. (2012b) (Fig. 1B) and in Anile et al. (2019). With respect to the methods reported in Anile et al. (2019), the 2018 survey was conducted from 30 May 2018 to 14 November 2018, with fewer cameras ($n = 76$ across seven line transects) and with a reduced trap-days effort ($n = 1,985$) due to the reduced availability of camera traps (Fig. 1B).

In the first photograph dated 26 May 2009 (Image

Editor: Angie Appel, Wild Cat Network, Bad Marienberg, Germany.

Date of publication: 17 February 2020 (online & print)

Citation: Anile, S., S. Devillard, C.K. Nielsen & M.L. Valvo (2020). Record of a 10-year old European Wildcat *Felis silvestris silvestris* Schreber, 1777 (Mammalia: Carnivora: Felidae) from Mt. Etna, Sicily, Italy. *Journal of Threatened Taxa* 12(2): 15272–15275. <https://doi.org/10.11609/jott.5484.12.2.15272-15275>

Copyright: © Anile et al. 2020. Creative Commons Attribution 4.0 International License. JoTT allows unrestricted use, reproduction, and distribution of this article in any medium by providing adequate credit to the author(s) and the source of publication.

Funding: This research was funded by Etna Park.

Competing interests: The authors declare no competing interests.

Acknowledgements: SA thanks Dr. Michele Leonardi and Dr. Rosa Spampinato (Etna Park officers) for making this study feasible. SA also thanks Dr. Luigi Piccinini and Maurizio Pennisi from the “Ripartizione Faunistico Venatoria di Catania” for providing the cameras used in this study. We also thank three anonymous reviewers for their useful comments which have improved a first version of this manuscript. Lastly, we all thank Marisa Mazzaglia, former president of Etna Park, for supporting this wildcat research without any hesitation.






Figure 1. The location of Mt. Etna (A) and European Wildcat distribution range. The inset (B) shows the study area with camera stations used during the 2009 and 2018 surveys.



Image 1. A & B—European Wildcat photographed at Mt. Etna on 26 May and 15 June 2009 | C—European Wildcat photographed at Mt. Etna on 10 June 2018. © Stefano Anile.

1A) this Wildcat was clearly identifiable by the absence of the typical black-tipped tail of the European Wildcat (Ragni & Possenti 1996); its tail showed only a clear white ring (Image 1B). In addition, the shape of the dorsal stripe aided to confirm its recapture. During the same survey, two additional photographs of this individual were obtained at two other camera stations. Collectively, three photographs at three neighbouring camera stations were recorded. During the camera trapping survey conducted in 2018, this individual Wildcat was recaptured on 10 June 2018 at one camera station (Image 1C), relatively near to where it was captured during the 2009 survey. The mean distance

from the other camera stations where it was previously recorded was 960m. The time between the first and last photograph was 3,302 days (9.04 years). Hence, the likely minimum age of this individual at the time of recapture must have been at least 10 years.

Camera trapping has greatly increased our scientific knowledge on many cat species worldwide. Indeed, many central topics for the proper management of Felidae have been investigated, such as population density estimation (Anile et al. 2014), habitat selection (Lesmeister et al. 2015; Anile et al. 2019), population dynamics (Karanth et al. 2006; Sharma et al. 2014; Duangchantrasiri et al. 2015; Majumder et al. 2017), and

adult sex ratio (Anile & Devillard 2018). Furthermore, recent years have seen an increased collaboration among researchers for sharing camera trapping data, and hence the investigation of ecological patterns at larger scales, i.e., across study areas (Steenweg et al. 2017; Davis et al. 2018; Khwaja et al. 2019).

The Wildcat population dwelling on Mt. Etna has been extensively ($n = 41$ individuals from 2010–2018) screened for detecting hybridization with Domestic Cats *Felis catus*, but no evidence of hybridization has been detected in this population (Mattucci et al. 2013; Anile et al. 2014, 2019). Hence, we consider more likely that a mutilation occurred at the end of the tail, which was also shorter than the normal size, ~30cm, that caused the loss of the black tip, rather than considering this anomaly in the typical Wildcat marking system pattern due to hybridization.

The Wildcat we recaptured after nine years was surely not a kitten at the time of the first capture, hence we think that 10 years is the minimum reasonable age estimation for this individual. This age estimate still lies at the lower range when compared to ages of captive Wildcats ranging from 12–16 years studied by Hartmann (2005).

Long-term camera trapping studies have been conducted on Tigers *Panthera tigris* and Ocelots *Leopardus pardalis*, however, the maximum ages attained were not reported (Karanth et al. 2006; Duangchantrasiri et al. 2015; Majumder et al. 2017; Satter et al. 2019). On the contrary, Harmsen et al. (2017) reported a maximum age of 14 and 13 years for male and female Jaguars *Panthera onca*, respectively. When comparing the maximum age we recorded with the few other longevity records of cat species from the wild (Hunter 2015), we note that our estimate is considerably high, but still within the range of those reported for the species, especially when considering the small body mass (3.7–4.9 kg; Johnson et al. 2017) of the European Wildcat.

The longevity of an individual European Wildcat might be influenced to some extent by local circumstances, e.g., absence of natural predators and widespread refuges as in our study area. In general, longevity is also shaped by ecological traits such as body mass (Healy et al. 2014), with larger species living longer than smaller ones. Long-term monitoring using camera traps can help understand patterns, which cannot be detected when using a small window in time, but this would require a more sustainable support from funding agencies as the costs involved with this kind of studies are certainly higher than surveys running over a shorter period.

References

- Anile, S. & S. Devillard (2015). Study design and body mass influence RAls from camera trap studies: evidence from the Felidae. *Animal Conservation* 19(1): 35–45. <https://doi.org/10.1111/acv.12214>
- Anile, S. & S. Devillard (2018). Camera-trapping provides insights into adult sex ratio variability in felids. *Mammal Review* 48(3): 168–179. <https://doi.org/10.1111/mam.12120>
- Anile, S., B. Ragni, E. Randi, F. Mattucci & F. Rovero (2014). Wildcat population density on the Etna volcano, Italy: a comparison of density estimation methods. *Journal of Zoology* 293(4): 252–261. <https://doi.org/10.1111/jzo.12141>
- Anile, S., C. Arrabito, M.V. Mazzamuto, D. Scornavacca & B. Ragni (2012a). A non-invasive monitoring on European Wildcat (*Felis silvestris silvestris* Schreber, 1777) in Sicily using hair trapping and camera trapping: does scented lure work? *Hystrix Italian Journal of Mammalogy* 23(2): 45–50. <https://doi.org/10.4404/hystrix-23.2-4657>
- Anile, S., C. Amico & B. Ragni (2012b). Population density estimation of the European Wildcat (*Felis silvestris*) in Sicily using camera trapping. *Wildlife Biology in Practice* 8(1): 1–12. <https://doi.org/10.2461/wbp.2012.8.1>
- Anile, S., L. Bizzarri & B. Ragni (2009). Camera trapping the European Wildcat (*Felis silvestris silvestris*) in Sicily (southern Italy): preliminary results. *Hystrix: Italian Journal of Mammalogy* 20(1): 55–60. <https://doi.org/10.4404/hystrix-20.1-4433>
- Anile, S., L. Bizzarri & B. Ragni (2010). Estimation of European Wildcat population size in Sicily (Italy) using camera trapping and capture–recapture analyses. *Italian Journal of Zoology* 77(2): 241–246. <https://doi.org/10.1080/11250000903419731>
- Anile, S., S. Devillard, B. Ragni, F. Rovero, F. Mattucci & M. Lo Valvo (2019). Habitat fragmentation and anthropogenic factors affect Wildcat (*Felis silvestris silvestris*) occupancy and detectability on Mt. Etna. *Wildlife Biology* 2019(1): 1–13. <https://doi.org/10.2981/wlb.00561>
- Brodie, J.F. (2009). Is research effort allocated efficiently for conservation? Felidae as a global case study. *Biodiversity Conservation* 18: 2927–2939. <https://doi.org/10.1007/s10531-009-9617-3>
- Cutler, R.G. (1979). Evolution of longevity in ungulates and carnivores. *Gerontology* 25: 69–86.
- Davis, C.L., L.N. Rich, Z.J. Farris, M.J. Kelly, M.S. Di Bitetti, Y. Di Blanco, S. Albanesi, M.S. Farhadinia, N. Gholikhani, S. Hamel, B.J. Harmsen, C. Wulfsch, M.D. Kane, Q. Martins, A.J. Murphy, R. Steenweg, S. Sunarto, A. Taktehrani, K. Thapa, J.M. Tucker, J. Whittington, F.A. Widodo, N.G. Yoccoz & D.A.W. Miller (2018). Ecological correlates of the spatial co-occurrence of sympatric mammalian carnivores worldwide. *Ecology Letters* 21(9): 1401–1412. <https://doi.org/10.1111/ele.13124>
- De Magalhaes, J.P. & J. Costa (2009). A database of vertebrate longevity records and their relation to other life-history traits. *Journal of Evolutionary Biology* 22: 1770–1774. <https://doi.org/10.1111/j.1420-9101.2009.01783.x>
- Duangchantrasiri, S., M. Umponjan, S. Simcharoen, A. Pattanavibool, S. Chaiwattana, S. Maneerat, N.S. Kumar, D. Devcharan Jathanna, A. Srivathsa & K.U. Karanth (2015). Dynamics of a low-density Tiger population in Southeast Asia in the context of improved law enforcement. *Biological Conservation* 30(3): 639–648. <https://doi.org/10.1111/cobi.12655>
- Gonzalez-Lagos, C., D. Sol & S.M. Reader (2010). Large-brained mammals live longer. *Journal of Evolutionary Biology* 23: 1064–1074. <https://doi.org/10.1111/j.1420-9101.2010.01976.x>
- Harmsen, B.J., R.J. Foster, E. Sanchez, C.E. Gutierrez-Gonzalez, S.C. Silver, L.E.T. Ostro, M.J. Kelly, E. Kay & H. Quigley (2017). Long term monitoring of Jaguars in the Cockscomb Basin Wildlife Sanctuary, Belize. implications for camera trap studies of carnivores. *PLoS ONE* 12(6): e0179505. <https://doi.org/10.1371/journal.pone.0179505>
- Hartmann, M. (2005). Reproduction and behaviour of European Wildcats in species-specific enclosures, 13pp. In: Herrmann, M.




- (ed). Symposium: Biology and conservation of the European Wildcat (*Felis silvestris silvestris*), Germany, January 21st–23rd 2005. Öko-Log, Parlow, Germany, 44pp.
- Healy, K., T. Guillerme, S. Finlay, A. Kane, S.B.A. Kelly, D. McClean, D.J. Kelly, I. Donohue, A.L. Jackson & N. Cooper (2014). Ecology and mode-of-life explain lifespan variation in birds and mammals. *Proceedings of the Royal Society B* 281: 20140298. <http://doi.org/10.1098/rspb.2014.0298>
- Hunter, L. (Ed.) (2015). *Wild Cats of the World*. Bloomsbury, London, New York, 240pp.
- Johnson, P.J., M.J. Noonan, A.C. Kitchener, L.A. Harrington, C. Newman & D.W. Macdonald (2017). Rensching cats and dogs: feeding ecology and fecundity trends explain variation in the allometry of sexual size dimorphism. *Royal Society Open Science* 4(6): 170453. <http://doi.org/10.1098/rsos.170453>
- Kamler, J.F. & D.W. Macdonald (2006). Longevity of a wild Bat-eared Fox. *South African Journal of Wildlife Research* 36(2): 199–200.
- Karanth, K.U., J.D. Nichols, N.S. Kumar & J.E. Hines (2006). Assessing Tiger population dynamics using photographic capture-recapture sampling. *Ecology* 87(11): 2925–2937. [https://doi.org/10.1890/0012-9658\(2006\)87\[2925:ATPDUP\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2006)87[2925:ATPDUP]2.0.CO;2)
- Khawaja, H., C. Buchan, O.R. Wearn, L. Bahaa-el-din, D. Bantlin, H. Bernard, R. Bitariho, T. Bohm, J. Borah, J. Brodie, W. Chutipong, B. du Preez, A. Ebang-Mbele, S. Edwards, E. Fairet, J.L. Frechette, A. Garside, L. Gibson, A. Giordano, G.V. Gopi, A. Granados, S. Gubbi, F. Harich, B. Haurez, R.W. Havmoller, O. Helmy, L.A. Isbell, K. Jenks, R. Kalle, A. Kamjing, D. Khamcha, C. Kiebou-Opepa, M. Kinnaird, C. Kruger, A. Laudisoit, A. Lynam, S.E. Macdonald, J. Mathai, J.M. Sienne, A. Meier, D. Mills, J. Mohd-Azlan, Y. Nakashima, H.C. Nash, D. Ngoprasert, A. Nguyen, T. O'Brien, D. Olson, C. Orbell, J. Poulsen, T. Ramesh, D. Reeder, R. Reyna, L.N. Rich, J. Rode-Margono, F. Rovero, D. Sheil, M.H. Shirley, K. Stratford, N. Sukumal, S. Suwanrat, N. Tantisuanuh, A. Tilker, T. Van Berkel, L.K. Van der Weyde, M. Varney, F. Weise, I. Wiesel, A. Wilting, S.T. Wong, C. Waterman & D.W.S. Challender (2019). Pangolins in global camera trap data: implications for ecological monitoring. *Global Ecology and Conservation* 20: e00769. <https://doi.org/10.1016/j.gecco.2019.e00769>
- Lesmeister, D.B., C.K. Nielsen, E.M. Schaubert & E.C. Hellgreen (2015). Spatial and temporal structure of a mesocarnivore guild in Midwestern North America. *Wildlife Monographs* 191: 1–61. <https://doi.org/10.1002/wmon.1015>
- Macdonald, D.W., A.J. Loveridge & K. Nowell (2010). Dramatis personae: an introduction to the wild felids. Pp. 3–58 in: Macdonald, D.W. & A.J. Loveridge (eds.). *The Biology and Conservation of Wild Felids*. Oxford University Press, Oxford, 783pp.
- Majumder, A., Q. Qureshi, K. Sankar & A. Kumar (2017). Long-term monitoring of a Bengal Tiger (*Panthera tigris tigris*) population in a human-dominated landscape of central India. *European Journal of Wildlife Research* 63(1): 1–11. <https://doi.org/10.1007/s10344-016-1070-5>
- Mattucci, F., R. Oliveira, L. Bizzarri, F. Vercillo, S. Anile, B. Ragni, L. Lapini, A. Sforzi, P.C. Alves, L.A. Lyons & E. Randi (2013). Genetic structure of Wildcat (*Felis silvestris*) populations in Italy. *Ecology and Evolution* 3: 2443–2458. <https://doi.org/10.1002/ece3.569>
- Ragni, B. & M. Possenti (1996). Variability of coat-colour and markings system in *Felis silvestris*. *Italian Journal of Zoology* 63: 285–292.
- Ricklefs, R.E. & C.D. Cadena (2007). Lifespan is unrelated to investment in reproduction in populations of mammals and birds in captivity. *Ecology Letters* 10: 867–875. <https://doi.org/10.1111/j.1461-0248.2007.01085.x>
- Satter, C.B., B.C. Augustine, B.J. Harmsen, R.J. Foster, E.E. Sanchez, C. Wultsch, M. Davis & M.J. Kelly (2019). Long-term monitoring of Ocelot densities in Belize. *Journal of Wildlife Management* 83(2): 283–294. <https://doi.org/10.1002/jwmg.21598>
- Sharma, K., R. Bayraksmith, L. Tumursukh, O. Johansson, P. Sevgier, T. McCarthy & C. Mishra (2014). Vigorous dynamics underlie a stable population of the endangered Snow Leopard *Panthera uncia* in Tost mountains, south Gobi, Mongolia. *PLoS ONE* 9(7): e101319. <https://doi.org/10.1371/journal.pone.0101319>
- Smith, J.E., K.D.S. Lehmann, T.M. Montgomery, E.D. Strauss & K.E. Holekamp (2017). Insights from long-term field studies of mammalian carnivores. *Journal of Mammalogy* 98(3): 631–641. <https://doi.org/10.1093/jmammal/gyw194>
- Steenweg, R., M. Hebblewhite, R. Kays, J. Ahumada, J.T. Fisher, C. Burton, S.E. Townsend, C. Carbone, J.M. Rowcliffe, J. Whittington, J. Brodie, J.A. Royle, A. Switalski, A.P. Clevenger, N. Heim & L.N. Rich (2017). Scaling-up camera traps: monitoring the planet's biodiversity with networks of remote sensors. *Frontiers in Ecology and Environment* 15(1): 26–34. <https://doi.org/10.1002/fee.1448>
- Tidieri, M., J.M. Gaillard, V. Berger, D.W.H. Muller, L.B. Lackey, O. Gimenez, M. Clauss & J.F. Lemaître (2016). Comparative analyses of longevity and senescence reveal variable survival benefits of living in zoos across mammals. *Scientific Reports* 6: 36361. <https://doi.org/10.1038/srep36361>
- UNESCO World Heritage Center (2019). United Nations World Heritage List: Mount Etna. Electronic version at <https://whc.unesco.org/en/list/1427> accessed on 6 January 2020.
- Wilkinson, G.S. & J.M. South (2002). Life history, ecology and longevity in bats. *Aging Cell* 1: 124–131





www.threatenedtaxa.org

PLATINUM
OPEN ACCESS



The Journal of Threatened Taxa (JoTT) is dedicated to building evidence for conservation globally by publishing peer-reviewed articles online every month at a reasonably rapid rate at www.threatenedtaxa.org. All articles published in JoTT are registered under [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/) unless otherwise mentioned. JoTT allows unrestricted use, reproduction, and distribution of articles in any medium by providing adequate credit to the author(s) and the source of publication.

ISSN 0974-7907 (Online) | ISSN 0974-7893 (Print)

February 2020 | Vol. 12 | No. 2 | Pages: 15219–15278

Date of Publication: 17 February 2020 (Online & Print)

DOI: 10.11609/jott.2020.12.2.15219-15278

Editorial

Foreword to the second special issue on small wild cats

– Angie Appel & Shomita Mukherjee, Pp. 15219–15220

Communications

Breaking barriers: Iberian Lynx *Lynx pardinus* Temminck, 1827 (Mammalia: Carnivora: Felidae) colonizing Olive groves

– Germán Garrote, José Francisco Bueno, Manuel Ruiz, Santiago de Lillo, José Manuel Martín, Manuel Moral & Miguel Ángel Simón, Pp. 15221–15228

Iberian Lynx *Lynx pardinus* Temminck, 1827 (Mammalia: Carnivora: Felidae) in central Spain: trophic niche of an isolated population

– Pedro Alfaya Herbello, Ariadna Invernón & Germán Alonso, Pp. 15229–15237

New records of the Flat-headed Cat *Prionailurus planiceps* (Vigors & Horsfield, 1827) (Mammalia: Carnivora: Felidae) in western Sarawak, Malaysia

– Jayasilan Mohd-Azlan & Syaza Jailan Thaqifah, Pp. 15238–15243

Temporal overlap between two sympatric carnivores in northwestern Peru and southwestern Ecuador

– Alvaro García-Olaechea & Cindy M. Hurtado, Pp. 15244–15250

Short Communications

First photographic record of Jungle Cat *Felis chaus* Schreber, 1777 (Mammalia: Carnivora: Felidae) in Haripur District, Pakistan

– Afzal Anjum, Angie Appel & Muhammad Kabir, Pp. 15251–15255

New record on Asiatic Golden Cat *Catopuma temminckii* Vigors & Horsfield, 1827 (Mammalia: Carnivora: Felidae): photographic evidence of its westernmost distribution in Gaurishankar Conservation Area, Nepal

– Narayan Prasad Koju, Bijay Bishyal, Bishnu Prasad Pandey, Shankar Thami, Man Kumar Dhamala & Satya Narayan Shah, Pp. 15256–15261

First photographic evidence of the Asiatic Golden Cat *Catopuma temminckii* Vigors & Horsfield, 1827 (Mammalia: Carnivora: Felidae) in Sakteng Wildlife Sanctuary, Bhutan

– Sonam Wangyel, Kumbu Dorji, Sonam Tobgay & Norbu Yangdon, Pp. 15262–15266

Jungle Cat *Felis chaus* Schreber, 1777 (Mammalia: Carnivora: Felidae) at high elevations in Annapurna Conservation Area, Nepal

– Bikram Shrestha, Naresh Subedi & Ram Chandra Kandel, Pp. 15267–15271

Notes

Record of a 10-year old European Wildcat *Felis silvestris silvestris* Schreber, 1777 (Mammalia: Carnivora: Felidae) from Mt. Etna, Sicily, Italy

– Stefano Anile, Sebastien Devillard, Clayton Kent Nielsen & Mario Lo Valvo, Pp. 15272–15275

Filling a gap in Andean Cat *Leopardus jacobita* (Cornalia, 1865) (Mammalia: Carnivora: Felidae) distribution range: new record in La Rioja province, Argentina

– Cintia Gisele Tellaeche, María de las Mercedes Guerisoli, Constanza Napolitano, Dante Luis Di Nucci & Juan Ignacio Reppucci, Pp. 15276–15278

Member



Publisher & Host

